



Bureau of Justice Statistics Bulletin

September 2007, NCJ 218915

HIV in Prisons, 2005

By Laura M. Maruschak
BJS Statistician

On December 31, 2005, correctional authorities reported that 20,888 State inmates (1.8%) and 1,592 Federal inmates (1.0%) were infected with the human immunodeficiency virus (HIV) or had confirmed AIDS. This was a total of 22,480 inmates, a slight decrease from 22,936 inmates in 2004.

Half of the HIV/AIDS cases were in the South, nearly a third in the Northeast, and about a tenth in both the Midwest and the West. The Northeast reported the highest percentage of HIV/AIDS cases based on its custody population (3.9%). At yearend 2005, three states — New York (4,440), Florida (3,396), and Texas (2,400) — housed nearly half (49%) of all HIV/AIDS cases in State prisons.

At yearend 2005, an estimated 18,953 males (1.8%) and 1,935 females (2.4%) in State prisons were HIV-infected or had confirmed AIDS. The number of cases for both males and females were down from 2004. At yearend 2005, five States reported no female HIV/AIDS cases: New Hampshire, North Dakota, West Virginia, Hawaii, and Montana. Among Federal inmates, 1,491 men (1.0%) and 101 women (0.9%) were HIV-infected or had confirmed AIDS.

An estimated 5,620 inmates had confirmed AIDS, down from 5,969 in 2004. Of the confirmed cases, 5,026 were in State prisons and 594 were in Federal prisons. Four States accounted for nearly half (47%) of all confirmed AIDS cases: Texas (894), New York (700), Florida (645), and Maryland (408). Maryland (1.8%) had the highest percentage of confirmed AIDS cases among its custody population, followed by Connecticut, New York, and Delaware (1.1% each). Five States (New Hampshire, Indiana, North Dakota, Idaho, and Wyoming) reported having no confirmed AIDS cases.

During 2005 an estimated 176 State inmates died from AIDS-related causes, down from 185 in 2004. For every 100,000 State inmates, 13 died from AIDS-related causes. AIDS-related deaths accounted for nearly 1 in 20 deaths reported in State prisons. New York reported the largest

Reported HIV/AIDS cases and AIDS-related deaths, 2004 and 2005

| | 2005 | | 2004 | |
|--|--------|-------------------------------|--------|-------------------------------|
| | Number | Percent of custody population | Number | Percent of custody population |
| U.S. total | 22,480 | 1.7% | 22,936 | 1.8% |
| Federal | 1,592 | 1.0% | 1,680 | 1.1% |
| State | 20,888 | 1.8% | 21,256 | 1.8% |
| Northeast | 6,456 | 3.9 | 6,646 | 4.1 |
| Midwest | 2,104 | 0.9 | 2,025 | 0.9 |
| South | 10,423 | 2.2 | 10,691 | 2.1 |
| West | 1,905 | 0.7 | 1,894 | 0.7 |
| Gender^a | | | | |
| Male | 20,444 | 1.7% | 19,294 | 1.8% |
| Female | 2,036 | 2.3 | 1,962 | 2.5 |
| Confirmed AIDS^a | 5,620 | 0.4% | 5,969 | 0.5% |
| AIDS-related deaths in prison^b | | | | |
| State | 176 | | 185 | |
| Federal | 27 | | 18 | |

^aIncludes estimates of the number inmates in States not reporting a breakdown of the total number of HIV/AIDS cases. Estimates are based on the most recent data available. See *Methodology*.

^bTotal number of deaths are based on a combination of data collected in the National Prisoners Statistics (NPS) and the Death in Custody Reporting Program (DCRP).

number of AIDS-related deaths (19), followed by Florida (17) and California (14). Relative to the number of inmates in custody, the Northeast reported the highest rate of AIDS-related deaths (28 per 100,000 inmates), followed by the South (13 per 100,000 inmates).

Among Federal inmates, 27 died from AIDS-related causes in 2005, up from 18 in 2004. For every 100,000 Federal inmates, 15 died from AIDS-related causes. AIDS-related deaths accounted for 7% of all deaths in Federal prisons.

State-level data and other detailed information are available in appendix tables on the BJS website at <<http://www.ojp.usdoj.gov/bjs/pub/hivp05.pdf>>.

HIV/AIDS cases in State and Federal prisons steadily declining since 1999

The Bureau of Justice Statistics (BJS) began collecting official data on HIV infection in State and Federal prisons in 1991 (see *Methodology*). At that time, 17,551 State and Federal inmates were HIV-infected or had confirmed AIDS (figure 1). This number steadily increased through yearend 1995, but fluctuated thereafter until reaching a high of 25,807 at yearend 1999. Since then, the number of HIV/AIDS cases has steadily declined, reaching 22,480 at yearend 2005.

New York accounted for more than three-quarters of the decline in HIV/AIDS cases between 1999 and 2005

Between 1999 and 2005, the number of inmates who were HIV-infected or had confirmed AIDS increased in 18 States and the Federal system, while the number decreased in 27 States (table 1). The remaining five States and the District of Columbia either reported no change or did not report data for 1999 or 2005.

The largest increases were reported in Florida (up 763), the Federal system (up 436), Georgia (up 196), and North Carolina (up 164). New York reported the largest decrease, with 2,560 fewer HIV-infected or confirmed AIDS inmates, followed by New Jersey (down 329), and California (down 321). New York alone accounted for more than three-quarters (83%) of the total reported decrease in comparable HIV/AIDS cases.¹

¹New York estimates the number of HIV/AIDS cases based on data obtained through blind sero-prevalence studies conducted biennially by the New York State Department of Health. See *Methodology* for details on the estimation procedures.




Figure 1

Table 1. Number of HIV/AIDS cases in State and Federal prisons at yearend, 1999 and 2005

| Jurisdiction | Total HIV/AIDS cases | | Change from 1999-2005 |
|-----------------------------------|----------------------|--------|-----------------------|
| | 2005 | 1999 | |
| U.S. total | | | |
| Comparable ^a | 22,346 | 25,438 | -3,092 |
| Reported | 22,480 | 25,807 | -3,327 |
| Federal | 1,592 | 1,156 | 436 |
| State | 20,888 | 24,651 | -3,763 |
| Northeast | 6,456 | 10,030 | -3,574 |
| Connecticut | 463 | 632 | -169 |
| Maine | 10 | 9 | 1 |
| Massachusetts | 221 | 346 | -125 |
| New Hampshire | 21 | 17 | 4 |
| New Jersey | 540 | 869 | -329 |
| New York | 4,440 | 7,000 | -2,560 |
| Pennsylvania | 692 | 939 | -247 |
| Rhode Island ^b | 58 | 203 | -145 |
| Vermont | 11 | 15 | -4 |
| Midwest | 2,104 | 2,171 | -67 |
| Illinois | 474 | 635 | -161 |
| Indiana | 134 | / | : |
| Iowa | 28 | 30 | -2 |
| Kansas | 34 | 41 | -7 |
| Michigan | 525 | 578 | -53 |
| Minnesota | 41 | 32 | 9 |
| Missouri | 301 | 290 | 11 |
| Nebraska | 19 | 20 | -1 |
| North Dakota | 2 | 2 | 0 |
| Ohio | 410 | 391 | 19 |
| South Dakota | 14 | 5 | 9 |
| Wisconsin | 122 | 147 | -25 |
| South | 10,423 | 10,243 | 180 |
| Alabama | 268 | 283 | -15 |
| Arkansas | 94 | 99 | -5 |
| Delaware | 124 | 170 | -46 |
| District of Columbia ^c | / | 359 | : |
| Florida | 3,396 | 2,633 | 763 |
| Georgia | 1,042 | 846 | 196 |
| Kentucky | 83 | 122 | -39 |
| Louisiana | 488 | 381 | 107 |
| Maryland | 671 | 820 | -149 |
| Mississippi | 302 | 192 | 110 |
| North Carolina | 718 | 554 | 164 |
| Oklahoma | 136 | 122 | 14 |
| South Carolina | 489 | 617 | -128 |
| Tennessee | 210 | 185 | 25 |
| Texas | 2,400 | 2,520 | -120 |
| Virginia | / | 330 | : |
| West Virginia | 2 | 10 | -8 |
| West | 1,905 | 2,207 | -302 |
| Alaska | / | 16 | : |
| Arizona | 152 | 144 | 8 |
| California | 1,249 | 1,570 | -321 |
| Colorado | 148 | 131 | 17 |
| Hawaii | 23 | 30 | -7 |
| Idaho | 26 | 15 | 11 |
| Montana | 6 | 10 | -4 |
| Nevada | 124 | 125 | -1 |
| New Mexico | 25 | 26 | -1 |
| Oregon | / | 23 | : |
| Utah | 38 | 34 | 4 |
| Washington | 107 | 75 | 32 |
| Wyoming | 7 | 8 | -1 |

Note: For detailed State-level data for 2005, see Appendix table 1 at <<http://www.ojp.usdoj.gov/bjs/pub/hivp05.pdf>>.

/Not reported.

:Not calculated.

^aThe comparable numbers for 1999 and 2005 include data from the Federal system, 46 States, and the District of Columbia.

^bIn 2005 Rhode Island changed their method for reporting HIV/AIDS cases. As a result, Rhode Island revised their HIV/AIDS counts for 2000-2004.

^cThe percentage change was not calculated for the District of Columbia because at yearend 2001 responsibility for housing sentenced felons was transferred to the Federal Bureau of Prisons.

Number of confirmed AIDS cases fluctuated between 1999 and 2005

Between 1999 and 2005, the estimated number of confirmed AIDS cases in State and Federal prisons fluctuated, reaching a high of 7,039 in 1999 and a low of 5,607 in 2002 (table 2). Overall, the estimated number of confirmed cases decreased by 1,419 cases between 1999 and 2005. Since 2000, the estimated confirmed AIDS cases as a percent of all HIV/AIDS cases have remained stable at about 25%.

The number of confirmed AIDS cases was estimated because not all States reported counts in every year during the 7-year period between 1999 and 2005. For States that reported counts, the number of confirmed AIDS cases ranged between 6,642 in 1999 and 4,862 in 2002.

A total of 31 States and the Federal system provided data in all 7 years (see comparable column in table 2). The pattern of change in confirmed AIDS cases was similar across the three measures — estimated, reported, and comparable confirmed AIDS cases. The number fluctuated but decreased overall during the period.

Difference between the rate of confirmed AIDS cases in prisons and the general population shrinking

In each year since 1991, the rate of confirmed AIDS has been higher among prison inmates than in the U.S. general population, but the gap has been narrowing. At yearend 2005, the estimated rate of confirmed AIDS in State and Federal prisons was more than 2½ times higher than in the general population (table 3). In 1999 it was nearly 5 times higher.

The shrinking difference between the rate of confirmed AIDS cases in prisons and the general population between 1999 and 2005 resulted from a decrease in the number of confirmed AIDS cases in prisons (down 20%) and an increase in cases (up 44%) among the general population.

HIV/AIDS cases among both male and female State inmates declining, but at a faster rate for females

Since 1999 the estimated number of both male and female inmates who had HIV infection or confirmed AIDS has decreased. Among men, HIV/AIDS dropped from 22,175 cases to 18,953 between 1999 and 2005 (table 4). For women, the number of cases dropped from 2,402 to 1,935. The percentage of male inmates who were HIV positive or had confirmed AIDS decreased from 2.2% to 1.8%; the percentage for female inmates fell from 3.4% to 2.4%.

While both groups experienced a decrease in the number of HIV/AIDS cases, females had a larger decrease in the percentage of HIV/AIDS cases. This can be explained by the faster growing female population in prisons and a larger decrease in the number of HIV/AIDS cases among women.

Table 2. Number of confirmed AIDS cases in State and Federal prisons, at yearend 1999-2005

| Yearend | Number of confirmed AIDS cases | | | Estimated confirmed AIDS cases as a percent of total HIV/AIDS cases |
|---------|--------------------------------|-----------------------|-------------------------|---|
| | Estimated ^a | Reported ^b | Comparable ^c | |
| 1999 | 7,039 | 6,642 | 5,973 | 27.3% |
| 2000 | 6,275 | 5,676 | 5,453 | 24.8 |
| 2001 | 6,240 | 5,708 | 5,399 | 25.9 |
| 2002 | 5,607 | 4,862 | 4,788 | 23.5 |
| 2003 | 5,892 | 5,175 | 5,095 | 25.0 |
| 2004 | 5,969 | 5,425 | 5,206 | 26.0 |
| 2005 | 5,620 | 5,422 | 5,011 | 25.0 |

Note: For State-level data for 2004 and 2005, see Appendix table 3 at <<http://www.ojp.usdoj.gov/bjs/pub/hivp05.pdf>>.

^aIn States that did not report confirmed AIDS cases, estimates were made by applying to the total the percentages by type of HIV infection from the most recent year for which data were available. For each year, estimates do not include data from States that did not report sufficient data on HIV/AIDS cases.

^bExcludes 6 States in 1999, 9 in 2000, 7 in 2001, 13 in 2002, 14 in 2003, 11 in 2004, and 9 in 2005.

^cThe Federal system and 31 States reported data for all 7 years shown in the table.

Table 3. Percent with confirmed AIDS among State and Federal prisoners and U.S. general population, 1999-2005

| Yearend | Percent of population estimated to have confirmed AIDS | | Percent of AIDS cases in prisons by percent in U.S. general population |
|---------|--|--------------------------|--|
| | State and Federal prisoners | U.S. general population* | |
| 1999 | 0.58% | 0.12% | 4.8 |
| 2000 | 0.51 | 0.13 | 3.9 |
| 2001 | 0.50 | 0.14 | 3.6 |
| 2002 | 0.45 | 0.14 | 3.2 |
| 2003 | 0.47 | 0.15 | 3.1 |
| 2004 | 0.46 | 0.15 | 3.1 |
| 2005 | 0.43 | 0.16 | 2.7 |

Note: See *Methodology* for data sources and calculations.

*Based on persons age 13 or older in 1999 and age 15 or older thereafter. Excludes confirmed AIDS cases in State and Federal prisons.

Table 4. HIV/AIDS cases in State and Federal prisons, by gender, 1999-2005

| Yearend | Male inmates in custody | | Female inmates in custody | |
|---------|-------------------------|-----------------------|---------------------------|-----------------------|
| | Estimated number | Percent of population | Estimated number | Percent of population |
| 1999 | 22,175 | 2.2% | 2,402 | 3.4% |
| 2000 | 21,889 | 2.2 | 2,467 | 3.6 |
| 2001 | 20,357 | 2.0 | 2,196 | 3.1 |
| 2002 | 20,265 | 1.9 | 2,153 | 2.9 |
| 2003 | 20,010 | 1.9 | 2,126 | 2.8 |
| 2004 | 19,294 | 1.8 | 1,962 | 2.5 |
| 2005 | 18,953 | 1.8 | 1,935 | 2.4 |

Note: To provide year-to-year comparisons, estimates were made for States that did not report data by gender. For each State, estimates were calculated by applying to the total the percentages by gender from the most recent year for which data were available. For State-level data for 2005, see Appendix table 2 at <<http://www.ojp.usdoj.gov/bjs/pub/hivp05.pdf>>.

Inmate survey data support decline in HIV infection in 1997 and 2004

Data reported by inmates in personal interviews, collected in the BJS Survey of Inmates in State Correctional Facilities, were consistent with the decline in HIV infection observed in National Prisoner Statistics (NPS-1) official records. Personal interview data are obtained every 5 to 6 years. The most recent data collections were in 1997 and 2004. In each year, inmates were asked if they had been tested for HIV at admission or had ever been tested and the results of the test. In 2004, 1.6% of tested State inmates reported being HIV positive, down from 2.2% in 1997.

Decreases found in the percent of HIV positive among male and female inmates and inmates ages 25 to 34

The inmate survey data provide a broad range of individual characteristics, including gender, race, age, current offense, and prior drug use, that can be used to describe trends. Despite apparent declines within each category, not all declines were statistically significant (see table footnote "a" for statistically significant items).

Overall, the percentage of State inmates reporting HIV infection in 2004 was down from 1997. Both male and female inmates were less likely to report being HIV positive in 2004 than in 1997. In 2004, 1.6% of male inmates, compared to 2.2% in 1997, were HIV positive. For female inmates, 1.9% in 1997, down from 3.4% in 2004, reported being HIV positive.

Among State inmates ages 25 to 34, there was a sharp decline in the number who reported being HIV positive between 1997 and 2004 (2.3% compared to 0.8%). Over the same time period, inmates in each of the other age categories were equally likely to report being HIV positive.

Decrease in HIV infection found among drug offenders and inmates who reported prior drug involvement

Drug offenders were less likely in 2004 to report being HIV positive than in 1997. In 2004, 1.8% of drug offenders reported being HIV positive, down from 2.9% in 1997. Between 1997 and 2004, significant decreases in the percentage of inmates who reported being HIV positive were found among inmates who ever used drugs, used drugs in the month before arrest, and used needles to inject drugs. Among inmates who reported

State inmates who were ever tested for HIV and reported results

| Characteristic | 2004 | | 1997 | |
|--|---------|----------------------|---------|----------------------|
| | Number | Percent HIV positive | Number | Percent HIV positive |
| All inmates | 967,200 | 1.6% | 790,100 | 2.2% |
| Gender | | | | |
| Male ^a | 897,700 | 1.6% | 734,300 | 2.2% |
| Female ^a | 69,500 | 1.9 | 55,800 | 3.4 |
| Race/Hispanic origin | | | | |
| White ^b | 336,700 | 1.0% | 257,900 | 1.4% |
| Black or African American ^b | 409,000 | 2.0 | 384,900 | 2.8 |
| Hispanic or Latino | 166,500 | 1.8 | 123,700 | 2.5 |
| Age | | | | |
| 24 or younger | 159,600 | 0.3% | 154,200 | 0.5% |
| 25-34 ^a | 328,900 | 0.8 | 310,200 | 2.3 |
| 35-44 | 300,600 | 2.5 | 232,800 | 3.1 |
| 45 or older | 178,000 | 2.6 | 92,200 | 2.7 |
| Current offense | | | | |
| Violent | 457,900 | 1.3% | 360,400 | 1.9% |
| Drug ^a | 201,800 | 1.8 | 164,300 | 2.9 |
| Property | 183,800 | 2.6 | 178,600 | 2.4 |
| Public-order | 110,700 | 0.9 | 77,000 | 1.9 |
| Prior drug use | | | | |
| Never | 155,400 | 1.3% | 123,000 | 1.7% |
| Ever ^a | 809,900 | 1.7 | 666,000 | 2.3 |
| In month before arrest ^a | 552,300 | 1.9 | 460,700 | 2.7 |
| Used needle to inject drugs ^a | 183,800 | 2.8 | 168,400 | 4.6 |
| Shared a needle | 77,900 | 5.1 | 74,400 | 7.7 |

Note: See Appendix table 7 for standard error estimates at <<http://www.ojp.usdoj.gov/bjs/pub/hivp05.pdf>>.

^aDifferences significant at the 95% confidence interval.

^bNot Hispanic or Latino.

ever using drugs, 1.7% reported being HIV positive in 2004, down from 2.3% in 1997. Inmates who used drugs in the month before arrest were less likely to report being HIV positive in 2004 (1.9%) than in 1997 (2.7%). Among inmates who said they had used a needle to inject drugs, 2.8% reported being HIV positive in 2004, down from 4.6% in 1997.

Number of AIDS-related deaths declined dramatically between 1995 and 2000

The number of State inmates who died of AIDS-related diseases, such as *Pneumocystis carinii* pneumonia and Kaposi's sarcoma,² peaked in 1995 at 1,010 deaths and declined through 2000 to 185 deaths. In 2001 the number of AIDS-related deaths reported in the NPS-1 increased to 256 and has steadily declined since (figure 2).

In 2001 BJS began collecting individual-level information about deaths in State prisons, under the Death in Custody Reporting Program (DCRP).³ The estimated number of AIDS-related deaths calculated using data from both the NPS-1 and the DCRP followed the same downward trend between 2001 and 2005 that was found using data from NPS-1 only. The estimated number of AIDS-related deaths was 176 in 2005, down from 311 in 2001.

Between 1995 and 2000, New York reported the largest decline in AIDS-related deaths (down 250), followed by Florida (down 102) and California (down 68). These three States combined accounted for more than half of the total decline (825) in the number of AIDS-related deaths. New Jersey (down 46) and Georgia (down 42) also had large declines.

²Center for Disease Control, 1993, Revised Classification System for HIV Infection and Expanded Surveillance Case Definition for AIDS Among Adolescents and Adults. *MMWR* 1992; 41(RR-17).

³In 2001 BJS began to estimate the number of AIDS-related deaths using data from the NPS-1 and the DCRP. The most complete data for each State's number of deaths were used to calculate the estimated number of AIDS-related deaths. See *Methodology*.

Trends in AIDS-related deaths, 1991-2005




Figure 2

Almost all of the decrease in AIDS-related deaths attributed to male inmates

Using the individual records collected under the DCRP, estimates of AIDS-related deaths by gender, age, and race were calculated by applying the percentages in DCRP in each year to the estimated total number of deaths. The number of AIDS-related deaths decreased across gender, age, race and Hispanic origin. Between 2001 and 2005, male inmates, inmates ages 35 to 44, and black non-Hispanic inmates accounted for most of the decrease (table 5).

In 2005, 166 male inmates died from AIDS-related causes, down from 295 in 2001. This decrease of 129 AIDS-related deaths among male inmates was almost all (96%) of the total decrease of 135 deaths among all inmates.

Table 5. Profile of inmates who died of AIDS-related causes in State prisons, 2001-2005

| Characteristic | Estimated number of AIDS-related deaths ^a | | | | | Rate of AIDS-related deaths per 100,000 inmates ^b | | | | |
|--|--|------|------|------|------|--|------|------|------|------|
| | 2005 | 2004 | 2003 | 2002 | 2001 | 2005 | 2004 | 2003 | 2002 | 2001 |
| State total | 176 | 185 | 268 | 283 | 311 | 14 | 15 | 22 | 24 | 26 |
| Gender | | | | | | | | | | |
| Male | 166 | 176 | 258 | 273 | 295 | 14 | 15 | 23 | 24 | 27 |
| Female | 10 | 9 | 10 | 10 | 16 | 12 | 11 | 12 | 13 | 21 |
| Age | | | | | | | | | | |
| 19 or younger | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 | 0 |
| 20-24 | 0 | 3 | 1 | 0 | 5 | 0 | 1 | 1 | 0 | 3 |
| 25-34 | 25 | 26 | 48 | 32 | 52 | 6 | 6 | 12 | 8 | 13 |
| 35-44 | 82 | 88 | 124 | 139 | 150 | 21 | 23 | 33 | 38 | 41 |
| 45-54 | 55 | 57 | 75 | 96 | 86 | 31 | 33 | 44 | 57 | 52 |
| 55 or older | 14 | 11 | 19 | 16 | 18 | 22 | 19 | 31 | 27 | 31 |
| Race/Hispanic origin | | | | | | | | | | |
| White ^c | 33 | 46 | 46 | 58 | 56 | 8 | 11 | 10 | 13 | 13 |
| Black or African American ^c | 120 | 122 | 186 | 188 | 208 | 24 | 24 | 34 | 35 | 39 |
| Hispanic or Latino | 21 | 11 | 32 | 35 | 46 | 9 | 5 | 15 | 17 | 22 |

Note: For State-level data for 2005, see appendix table 4 at <<http://www.ojp.usdoj.gov/bjs/pub/hivp05.pdf>>.

^aEstimates of the number of AIDS-related deaths by gender, age, and race/Hispanic origin were made by applying the percentages based on DCRP data to the estimated total number of AIDS-related deaths.

^bTo calculate the rate, the number of State prisoners by age was first estimated by applying the age distribution reported in the 2004 Survey of Inmates in State Correctional Facilities to the 2001-2005 midyear custody counts in NPS-1.

^cNot Hispanic or Latino.

Between 2001 and 2005, inmates ages 35 to 44 accounted for half of the decrease (down 68) in the total number of AIDS-related deaths. AIDS-related deaths of black non-Hispanic inmates accounted for 65% of the total decline of 88 AIDS-related deaths reported between 2001 and 2005 (down from 208 to 120).

AIDS-related deaths as a percent of all deaths decreasing more rapidly in State prison than in the general population

AIDS-related deaths as a percent of total deaths in State prisons also decreased significantly between 1995 and 2000, from 34.2% to 5.3% (table 6). Over a slightly different time period (1995 to 2004, the most recent year for which data are available), AIDS-related deaths among the general population ages 15 to 54 decreased 68%, from 44,233 to 14,315. AIDS-related deaths as a percent of all deaths in the general population declined from 12.9% to 4.3% during the period.

Between 2001 and 2004, the rate of AIDS-related deaths among prison inmates as a percent of all deaths in the prison population was nearly cut in half (from 10.3% to 5.6%). However, the rate in the general population remained stable at about 4.3%.

Rate of AIDS-related deaths declining in prisons, stabilizing in the general population

Between 1995 and 2001, the rate of AIDS-related deaths declined from 100 deaths per 100,000 inmates to 25 per 100,000 (table 7). In the general population the rate dropped from 29 per 100,000 to 9 per 100,000. After 2001, while the rate of AIDS-related deaths in the State prison population continued to decline, from 25 to 13 per 100,000 inmates, the rate in the general population ages 15 to 54 remained stable between 8 and 9 deaths per 100,000 persons.

Older persons in State prisons had a higher rate of AIDS-related deaths than those in the general population

Older inmates were more likely to die from AIDS-related causes than persons in the same age group in the general population. In 2005, inmates ages 35 to 44 were almost twice as likely to die from AIDS-related causes (21 per 100,000 State inmates) than their counterparts in the general population (12 per 100,000 persons) (table 8). Among persons ages 45 to 54, State inmates were more than twice as likely as those in the general population to die from AIDS-related causes (31 per 100,000 compared to 14 per 100,000).

Table 6. Percent of AIDS-related deaths among all deaths in State prisons and U.S. general population

| Yearend | Percent of deaths | | Percent of deaths in State prisons by percent in U.S. general population, ages 15-54 |
|---------|----------------------------|--|--|
| | State prisons ^a | U.S. general population, ages 15-54 ^b | |
| 1995 | 34.2% | 12.9% | 2.6 |
| 2001 | 10.3 | 4.3 | 2.4 |
| 2002 | 9.1 | 4.1 | 2.2 |
| 2003 | 8.0 | 4.2 | 1.9 |
| 2004 | 5.6 | 4.3 | 1.3 |
| 2005 | 5.3 | ... | : |

... Not available.

:Not calculated.

^aPercentages were based on the number of inmate deaths, excluding those in jurisdictions not reporting AIDS-related deaths.

^bExcludes deaths reported in State prisons. See *Methodology* for data sources.

Table 7. Rate of AIDS-related deaths in State prisons and U.S. general population

| Yearend | Rate per 100,000 persons | | Rate of deaths in State prisons by rate in U.S. general population, ages 15-54 ^b |
|---------|--------------------------|--|---|
| | State prisons | U.S. general population, ages 15-54 ^a | |
| 1995 | 100 | 29 | 3.5 |
| 2001 | 25 | 9 | 2.9 |
| 2002 | 22 | 9 | 2.6 |
| 2003 | 21 | 9 | 2.4 |
| 2004 | 14 | 9 | 1.7 |
| 2005 | 13 | 8 | 1.7 |

^aExcludes deaths reported in State prisons. See *Methodology* for data sources.

^bCalculation based on unrounded rates.

Table 8. Rate of AIDS-related deaths per 100,000 persons in State prisons and U.S. general population, by age, 2001-2005

| Age | Rate of AIDS-related deaths per 100,000 persons, by age ^a | | | | Percent of population | | |
|-------|--|------|--------------------------------------|------|-----------------------|------|-------------------------|
| | State prisons | | U.S. general population ^b | | State prisons | | U.S. general population |
| | 2005 | 2001 | 2005 | 2001 | 2004 | 2005 | 2001 |
| 20-24 | 0 | 3 | 1 | 1 | 15.5% | 7.1% | 6.9% |
| 25-34 | 6 | 13 | 4 | 6 | 33.1 | 13.5 | 13.8 |
| 35-44 | 21 | 42 | 12 | 15 | 30.5 | 14.7 | 15.8 |
| 45-54 | 31 | 52 | 14 | 13 | 14.1 | 14.3 | 13.8 |

Note: Data on AIDS-related deaths by age were not collected in prisons prior to 2001.

^aTo calculate the rate, the number of State prisoners by age was first estimated by applying the age distribution reported in the 2004 Survey of Inmates in State Correctional Facilities to the 2005 and 2001 mid-year custody counts in NPS-1.

^bExcludes deaths reported in State prisons. See *Methodology* for data sources.

Methodology

National Prisoner Statistics

The National Prisoner Statistics collection (NPS-1), which primarily measures prison population movement, began in 1926. The NPS-1 includes yearend counts of prisoners by jurisdiction, gender, race, Hispanic origin, and admissions and releases during the year. The series consists of reports from the departments of corrections in the 50 States and the Federal Bureau of Prisons. In 1991 BJS began collecting data on HIV/AIDS in prisons in NPS-1. BJS respondents have indicated the circumstances under which inmates are tested for HIV and have provided the number of HIV-infected inmates in their custody.

Surveys of Inmates in State Correctional Facilities

The Survey of Inmates in State Correctional Facilities provides nationally representative data on State prison inmates. The 1997 and 2004 surveys used the same questionnaire and a stratified two-stage sample design. Facilities were selected in the first stage and inmates were selected for interviews in the second stage. Detailed descriptions of each survey's methodology and links to other reports or findings are available on the BJS Website <<http://www.ojp.usdoj.gov/bjs/abstract/sicf04.htm>>.

Accuracy of the survey estimates

The accuracy of the estimates in this report depends on two types of error: sampling and nonsampling. Sampling error is the variation that may occur by chance because a sample rather than a complete enumeration of the population was conducted. Nonsampling error can be attributed to many sources, such as nonresponses, differences in the interpretation of questions among inmates, recall difficulties, and processing errors. In any survey the full extent of the nonsampling error is never known. The sampling error, as measured by an estimated standard error, varies by the size of the estimate and the size of the base population.

Testing for differences in the proportions of inmates reporting that they were HIV positive within subgroups in the 1997 and 2004 surveys compared a calculated z-value against the critical value of 1.645 (the 95% level for a one-tail test). Given the overall declines in HIV cases reported in the administrative data, a one-tail test showed that the difference in proportions was less than zero. The z-value was calculated by taking the ratio of the estimated difference in proportions to the square root of the pooled variances of the two samples. (See standard errors in appendix table 7 at <<http://www.ojp.usdoj.gov/pub/hiv05.pdf>>.)

Deaths in Custody Reporting Program

To implement the Death in Custody Reporting Act of 2000 (PL 106-297), BJS developed four quarterly data collections of death records from local jails (began in 2000), State prisons (2001), State juvenile correctional agencies (2002),

and State and local law enforcement agencies (2003). Records include data on the deceased's characteristics (such as age, gender, and race/Hispanic origin), criminal background (such as legal status, offenses, and time in custody), and the death itself (such as cause, time, location, and medical conditions and treatment).

Estimation of HIV/AIDS cases in New York State

New York estimates the number of HIV/AIDS cases based on data from blind sero-prevalence studies conducted biennially by the New York State Department of Health. Blood samples are taken from all inmates entering New York State prisons. Every other year an extra sample from 1,000 sequential admissions at reception in four reception centers is tested for various diseases, including HIV. The percentage with HIV infection is applied to the total inmate population, then adjusted for length of stay and data from other studies. Projections for interim years are made without blind studies.

AIDS in the U.S. resident population

The number of persons with confirmed AIDS in the U.S. general population (age 13 and older) was derived from the Centers for Disease Control and Prevention (CDC), HIV/AIDS Surveillance Report, yearend editions 1995-2005. For each year the number of active AIDS cases in the United States was calculated by taking the —

- cumulative number of total AIDS cases for persons age 13 or older at yearend (from the HIV/AIDS Surveillance Report), minus estimated confirmed AIDS cases in State and Federal prisons, and subtracting the
- cumulative number of AIDS deaths for people age 15 or older at yearend, minus estimated number of AIDS-related deaths in State and Federal prisons.

The rate of confirmed AIDS cases in the U.S. general population was calculated by —

- dividing the annual total number of individuals with AIDS by
- the estimated U.S. general population (age 13 or older before 2000; age 15 or older since 2000) minus the State and Federal custody population.

AIDS-related deaths in the United States

The number of AIDS-related deaths for persons ages 15 to 54 was based on the CDC, HIV/AIDS Surveillance Report, yearend editions. Deaths in the U.S. population for persons ages 15 to 54 were taken from the CDC, Monthly Vital Statistics Report, Vol. 42, No. 2(S); Vol. 43, No. 12; Vol. 43, No. 6(S); Vol. 45, No. 3(S); and Vol. 45, No. 11(S). Also deaths were taken from the CDC, National Vital Statistics Report, Vol. 47, No. 9; Vol. 47, No. 25; Vol. 48, No. 11; Vol. 49, No. 8; Vol. 50, No. 15; Vol. 53, No. 15; and Vol. 54, No. 19.



Washington, DC 20531

Official Business
Penalty for Private Use \$300

For 2001 to 2005, AIDS-related deaths were calculated as a percent of all deaths among persons ages 15 to 54 in the U.S. general population. The difference of the national estimate of AIDS deaths of persons ages 15 to 54 minus AIDS-related deaths of persons ages 15 to 54 in State prison was divided by the national mortality estimates of persons ages 15 to 54 minus total deaths in State prisons.

For 2001 through 2005, the rates of AIDS-related deaths in the general population were calculated by taking the difference of the national estimate of AIDS-related deaths for persons ages 15 to 54 minus AIDS-related deaths for those ages 15 to 54 in State prisons and dividing it by the U.S. general population estimate minus the State prison population ages 15 to 54.

Because data on AIDS-related deaths by age in State prisons were not collected prior to 2001, the total number of AIDS-related deaths in State prison were subtracted from the national estimate of AIDS-related deaths for the 1995 rate calculations.

The Bureau of Justice Statistics is the statistical agency of the U.S. Department of Justice. Jeffrey L. Sedgwick is the director.

BJS Bulletins present the first release of findings from permanent data collection programs.

This Bulletin was written by Laura M. Maruschak. Margaret Noonan and Heather Couture verified the report. Theresa M. Reitz carried out the data collection and processing under the supervision of Charlene M. Sebold, Governments Division, Census Bureau, U.S. Department of Commerce. Duane H. Cavanaugh and Diron J. Gaskins provided technical assistance. Tina Dorsey and Carolyn C. Williams edited the report and Jayne Robinson prepared it for final printing, under the supervision of Doris J. James.

September 2007, NCJ 218915

Office of Justice Programs

Innovation • Partnerships • Safer Neighborhoods
<http://www.ojp.usdoj.gov>

This report in portable document format and in ASCII and its related statistical data and tables are available at the BJS World Wide Web Internet site: <<http://www.ojp.usdoj.gov/bjs/abstract/hivp05.htm>>.

Appendix table 1. Inmates in custody of State or Federal prison authorities and reported to be positive for the human immunodeficiency virus (HIV) or to have confirmed AIDS, 2003-05

| Jurisdiction | Total HIV/AIDS cases ^a | | | HIV/AIDS cases as a percent of total custody population ^b | | |
|-----------------------------------|-----------------------------------|--------|--------|--|------|------|
| | 2005 | 2004 | 2003 | 2005 | 2004 | 2003 |
| U.S. total | | | | | | |
| Comparable reporting ^c | 22,263 | 22,560 | 23,247 | | | |
| Reported ^d | 22,480 | 22,936 | 23,608 | 1.7% | 1.8% | 1.9% |
| Federal | 1,592 | 1,680 | 1,631 | 1.0% | 1.1% | 1.1% |
| State | 20,888 | 21,256 | 21,977 | 1.8 | 1.8 | 2.0 |
| Northeast | 6,456 | 6,646 | 7,339 | 3.9% | 4.1% | 4.4% |
| Connecticut | 463 | 477 | 572 | 2.6 | 2.7 | 3.2 |
| Maine | 10 | 11 | 10 | 0.5 | 0.6 | 0.5 |
| Massachusetts | 221 | 215 | 230 | 2.1 | 2.2 | 2.3 |
| New Hampshire | 21 | 31 | 21 | 0.9 | 1.3 | 0.9 |
| New Jersey | 540 | 655 | 657 | 2.3 | 2.9 | 2.8 |
| New York | 4,440 | 4,500 | 5,000 | 7.0 | 7.0 | 7.6 |
| Pennsylvania | 692 | 693 | 775 | 1.7 | 1.7 | 1.9 |
| Rhode Island | 58 | 51 | 65 | 1.7 | 1.6 | 1.8 |
| Vermont | 11 | 13 | 9 | 0.7 | 0.9 | 0.6 |
| Midwest | 2,104 | 2,025 | 2,087 | 0.9% | 0.9% | 1.0% |
| Illinois | 474 | 488 | 551 | 1.1 | 1.1 | 1.3 |
| Indiana | 134 | / | / | 0.6 | / | / |
| Iowa | 28 | 32 | 31 | 0.3 | 0.4 | 0.4 |
| Kansas | 34 | 41 | 32 | 0.4 | 0.5 | 0.3 |
| Michigan | 525 | 575 | 579 | 1.1 | 1.2 | 1.2 |
| Minnesota | 41 | 44 | 40 | 0.5 | 0.6 | 0.5 |
| Missouri | 301 | 294 | 268 | 1.0 | 1.0 | 0.9 |
| Nebraska | 19 | 20 | 21 | 0.4 | 0.5 | 0.5 |
| North Dakota | 2 | 6 | 2 | 0.2 | 0.5 | 0.2 |
| Ohio | 410 | 387 | 416 | 1.0 | 0.9 | 1.0 |
| South Dakota | 14 | 9 | 11 | 0.4 | 0.3 | 0.4 |
| Wisconsin | 122 | 129 | 136 | 0.6 | 0.6 | 0.7 |
| South | 10,423 | 10,691 | 10,740 | 2.2% | 2.1% | 2.2% |
| Alabama | 268 | 270 | 270 | 1.1 | 1.1 | 1.0 |
| Arkansas | 94 | 102 | 94 | 0.8 | 0.8 | 0.7 |
| Delaware | 124 | 149 | 105 | 1.8 | 2.2 | 1.6 |
| Florida | 3,396 | 3,250 | 3,112 | 3.9 | 3.9 | 3.9 |
| Georgia | 1,042 | 1,109 | 1,095 | 2.1 | 2.2 | 2.3 |
| Kentucky | 83 | 74 | / | 0.7 | 0.6 | / |
| Louisiana | 488 | 487 | 511 | 2.5 | 2.5 | 2.6 |
| Maryland | 671 | 792 | 988 | 3.0 | 3.4 | 4.2 |
| Mississippi | 302 | 254 | 200 | 2.7 | 2.2 | 1.6 |
| North Carolina | 718 | 647 | 640 | / | 1.8 | 1.9 |
| Oklahoma | 136 | 133 | 162 | 0.8 | 0.8 | 1.0 |
| South Carolina | 489 | 489 | 520 | 2.2 | 2.2 | 2.3 |
| Tennessee | 210 | 215 | 208 | 1.5 | 1.5 | 1.5 |
| Texas | 2,400 | 2,405 | 2,460 | 1.7 | 1.7 | 1.8 |
| Virginia | / | 302 | 361 | / | 1.0 | 1.2 |
| West Virginia | 2 | 13 | 14 | 0.0 | 0.3 | 0.4 |
| West | 1,905 | 1,894 | 1,811 | 0.7% | 0.7% | 0.7% |
| Alaska | / | / | / | / | / | / |
| Arizona | 152 | 155 | 123 | 0.5 | 0.5 | 0.4 |
| California ^e | 1,249 | 1,212 | 1,196 | 0.7 | 0.7 | 0.7 |
| Colorado | 148 | 185 | 162 | 0.9 | 1.1 | 1.0 |
| Hawaii | 23 | 15 | 27 | 0.6 | 0.4 | 0.7 |
| Idaho | 26 | 33 | 26 | 0.6 | 0.7 | 0.6 |
| Montana | 6 | 5 | 4 | 0.3 | 0.2 | 0.2 |
| Nevada | 124 | 116 | 107 | 1.1 | 1.1 | 1.0 |
| New Mexico | 25 | 25 | 29 | 0.4 | 0.4 | 0.5 |
| Oregon | / | / | / | / | / | / |
| Utah | 38 | 39 | 37 | 0.8 | 0.8 | 0.8 |
| Washington | 107 | 102 | 93 | 0.6 | 0.6 | 0.6 |
| Wyoming | 7 | 7 | 7 | 0.6 | 0.6 | 0.6 |

/Not reported.

^aCounts published in previous reports may have been revised.

^bPercentages are based on custody counts, except for New Mexico for which percentages are based on its yearend jurisdiction count.

^cExcludes data from Indiana, Kentucky, Virginia, Alaska, and Oregon for all 3 years due to incomplete reporting.

^dExclude inmates in jurisdictions that did not report data.

^eThe 2005 number of HIV/AIDS cases in California was based on the count on 2/3/06.

Appendix table 2. Inmates in custody of State and Federal prison authorities reported to be positive for the human immunodeficiency virus (HIV) or to have confirmed AIDS, by gender, 2005

| Jurisdiction | Male HIV/AIDS cases | | Female HIV/AIDS cases | |
|--------------------------|---------------------|-----------------------|-----------------------|-----------------------|
| | Number | Percent of population | Number | Percent of population |
| U.S. total | | | | |
| Estimated ^{a,b} | 20,444 | | 2,036 | |
| Reported | 20,320 | 1.7% | 2,024 | 2.3% |
| Federal | 1,491 | 1.0% | 101 | 0.9% |
| State | 18,829 | 1.8 | 1,923 | 2.5 |
| Northeast | 5,839 | 3.7% | 617 | 7.2% |
| Connecticut | 411 | 2.5 | 52 | 4.0 |
| Maine | 9 | 0.5 | 1 | 0.8 |
| Massachusetts | 195 | 2.0 | 26 | 3.3 |
| New Hampshire | 21 | 0.9 | 0 | 0 |
| New Jersey | 463 | 2.1 | 77 | 6.8 |
| New York | 4,040 | 6.7 | 400 | 14.1 |
| Pennsylvania | 636 | 1.6 | 56 | 2.9 |
| Rhode Island | 54 | 1.7 | 4 | 2.1 |
| Vermont | 10 | 0.7 | 1 | 0.7 |
| Midwest | 1,950 | 0.8% | 154 | 0.9% |
| Illinois | 430 | 1.0 | 44 | 1.6 |
| Indiana | 117 | 0.6 | 17 | 0.9 |
| Iowa | 24 | 0.3 | 4 | 0.5 |
| Kansas | 33 | 0.4 | 1 | 0.1 |
| Michigan | 487 | 1.0 | 38 | 1.8 |
| Minnesota | 40 | 0.5 | 1 | 0.2 |
| Missouri | 287 | 1.0 | 14 | 0.6 |
| Nebraska | 18 | 0.5 | 1 | 0.2 |
| North Dakota | 2 | 0.2 | 0 | 0 |
| Ohio | 380 | 1.0 | 30 | 0.9 |
| South Dakota | 13 | 0.4 | 1 | 0.3 |
| Wisconsin | 119 | 0.6 | 3 | 0.2 |
| South | 9,306 | 2.2% | 981 | 3.1% |
| Alabama | 256 | 1.1 | 12 | 1.0 |
| Arkansas | 85 | 0.7 | 9 | 1.2 |
| Delaware | 103 | 1.6 | 21 | 3.9 |
| Florida | 3,054 | 3.8 | 342 | 5.7 |
| Georgia | 949 | 2.1 | 93 | 3.2 |
| Kentucky | 70 | 0.6 | 13 | 1.4 |
| Louisiana | 455 | 2.5 | 33 | 3.2 |
| Maryland | 597 | 2.8 | 74 | 6.9 |
| Mississippi | 262 | 2.7 | 40 | 2.5 |
| North Carolina | 654 | 1.9 | 64 | 2.4 |
| Oklahoma | / | / | / | / |
| South Carolina | 452 | 2.2 | 37 | 2.5 |
| Tennessee | 203 | 1.6 | 7 | 0.6 |
| Texas | 2,164 | 1.7 | 236 | 2.3 |
| Virginia | / | / | / | / |
| West Virginia | 2 | 0.1 | 0 | 0 |
| West | 1,734 | 0.7% | 171 | 0.8% |
| Alaska | / | / | / | / |
| Arizona | 138 | 0.5 | 14 | 0.5 |
| California | 1,149 | 0.7 | 100 | 0.9 |
| Colorado | 135 | 0.9 | 13 | 0.7 |
| Hawaii | 23 | 0.7 | 0 | 0 |
| Idaho | 23 | 0.6 | 3 | 0.6 |
| Montana | 6 | 0.4 | 0 | 0 |
| Nevada | 99 | 0.9 | 25 | 2.7 |
| New Mexico | 23 | 0.4 | 2 | 0.3 |
| Oregon | / | / | / | / |
| Utah | 32 | 0.7 | 6 | 1.3 |
| Washington | 100 | 0.7 | 7 | 0.5 |
| Wyoming | 6 | 0.5 | 1 | 0.9 |

/Not reported.

^aIncludes estimates of the number of inmates with HIV/AIDS by gender for Oklahoma.

^bIn States that did not report HIV/AIDS cases by gender, estimates were made by applying to the total the percentages by gender from the most recent year for which data were available. For each year, estimates do not include data from States that did not report sufficient data on HIV/AIDS cases.

Appendix table 3. Inmates in custody of State or Federal prison authorities and reported to have confirmed AIDS, yearend 2004 and 2005

| Jurisdiction | Confirmed AIDS cases ^a | | | |
|------------------------|-----------------------------------|-------|--|------|
| | Number | | Percent of custody population ^b | |
| | 2005 | 2004 | 2005 | 2004 |
| U.S. total | | | | |
| Estimated ^c | 5,620 | 5,969 | | |
| Reported | 5,422 | 5,425 | 0.5% | 0.5% |
| Federal | 594 | 641 | 0.4% | 0.4% |
| State | 4,828 | 4,784 | 0.5 | 0.5 |
| Northeast | 1,322 | 1,500 | 0.8% | 0.9% |
| Connecticut | 200 | 213 | 1.1 | 1.2 |
| Maine | 1 | 0 | -- | 0 |
| Massachusetts | 101 | 83 | 1.0 | 0.8 |
| New Hampshire | 0 | 6 | 0 | 0.2 |
| New Jersey | 118 | 202 | 0.5 | 0.9 |
| New York | 700 | 800 | 1.1 | 1.2 |
| Pennsylvania | 197 | 192 | 0.5 | 0.5 |
| Rhode Island | / | / | / | / |
| Vermont | 5 | 4 | 0.3 | 0.3 |
| Midwest | 497 | 389 | 0.2% | 0.2% |
| Illinois | 96 | 167 | 0.2 | 0.4 |
| Indiana | 0 | / | 0 | / |
| Iowa | 13 | 16 | 0.1 | 0.2 |
| Kansas | 4 | 6 | -- | 0.1 |
| Michigan | 223 | / | 0.5 | / |
| Minnesota | 21 | 18 | 0.3 | 0.2 |
| Missouri | / | 4 | / | -- |
| Nebraska | 8 | 8 | 0.2 | 0.2 |
| North Dakota | 0 | 1 | 0 | 0.1 |
| Ohio | 89 | 141 | 0.2 | 0.3 |
| South Dakota | 3 | 1 | 0.1 | -- |
| Wisconsin | 40 | 27 | 0.2 | 0.1 |
| South | 2,670 | 2,565 | 0.7% | 0.7% |
| Alabama | 29 | 133 | 0.1 | 0.5 |
| Arkansas | 24 | 13 | 0.2 | 0.1 |
| Delaware | 74 | 61 | 1.1 | 0.9 |
| Florida | 645 | 721 | 0.7 | 0.9 |
| Georgia | / | / | / | / |
| Kentucky | 12 | 53 | 0.1 | 0.5 |
| Louisiana | / | / | / | / |
| Maryland | 408 | 204 | 1.8 | 0.9 |
| Mississippi | 46 | 51 | 0.4 | 0.4 |
| North Carolina | 295 | 249 | 0.8 | 0.7 |
| Oklahoma | / | / | / | / |
| South Carolina | 196 | 212 | 0.9 | 0.9 |
| Tennessee | 45 | 66 | 0.3 | 0.5 |
| Texas | 894 | 801 | 0.6 | 0.6 |
| Virginia | / | / | / | / |
| West Virginia | 2 | 1 | -- | -- |
| West | 339 | 330 | 0.1% | 0.1% |
| Alaska | / | / | / | / |
| Arizona | 11 | / | -- | / |
| California | 196 | 191 | 0.1 | 0.1 |
| Colorado | 33 | 38 | 0.2 | 0.2 |
| Hawaii | 2 | 2 | 0.1 | 0.1 |
| Idaho | 0 | / | 0 | / |
| Montana | 1 | 0 | 0.1 | 0 |
| Nevada | / | 19 | / | 0.2 |
| New Mexico | 4 | 4 | 0.1 | 0.1 |
| Oregon | / | / | / | / |
| Utah | 22 | 11 | 0.4 | 0.2 |
| Washington | 70 | 65 | 0.4 | 0.4 |
| Wyoming | 0 | 0 | 0 | 0 |

/Not reported.

--Less than .05%.

^aCounts published in previous reports have been revised.

^bPercents exclude inmates in jurisdictions that did not report type of HIV/AIDS infection.

^cIncludes estimates of the number of inmates with confirmed AIDS for Georgia, Louisiana, and Oklahoma in 2004 and 2005; for Arizona, Idaho, Indiana, Michigan, and Virginia in 2004; and for Missouri and Nevada in 2005. Also see table 2, footnote a.

Appendix table 4. Deaths of State prisoners, 2005

| Jurisdiction | Prisoner deaths reported | | | | Total estimated deaths ^b | | | |
|------------------|--------------------------|--------------|----------------------|--------------|-------------------------------------|--------------------------|--------------|--------------------------|
| | In NPS-1 | | In DCRP ^a | | All causes | | AIDS-related | |
| | Total | AIDS-related | Total | AIDS-related | Number | Rate per 100,000 inmates | Number | Rate per 100,000 inmates |
| Total | 3,308 | 121 | 3,179 | 172 | 3,316 | 249 | 176 | 13 |
| Northeast | 477 | 41 | 474 | 48 | 477 | 276 | 48 | 28 |
| Connecticut | 41 | 9 | 40 | 11 | 41 | 208 | 11 | 56 |
| Maine | 1 | 0 | 0 | 0 | 1 | 48 | 0 | 0 |
| Massachusetts | 37 | 0 | 37 | 0 | 37 | 353 | 0 | 0 |
| New Hampshire | 8 | 0 | 8 | 0 | 8 | 312 | 0 | 0 |
| New Jersey | 61 | 5 | 61 | 6 | 61 | 217 | 6 | 21 |
| New York | 172 | 17 | 171 | 19 | 172 | 273 | 19 | 30 |
| Pennsylvania | 149 | 10 | 149 | 11 | 149 | 359 | 11 | 26 |
| Rhode Island | 2 | 0 | 2 | 0 | 2 | 55 | 0 | 0 |
| Vermont | 6 | 0 | 6 | 1 | 6 | 304 | 1 | 51 |
| Midwest | 557 | 10 | 542 | 24 | 557 | 220 | 24 | 9 |
| Illinois | 74 | 5 | 73 | 6 | 74 | 166 | 6 | 13 |
| Indiana | 50 | 2 | 45 | 2 | 50 | 204 | 2 | 8 |
| Iowa | 19 | 0 | 19 | 0 | 19 | 221 | 0 | 0 |
| Kansas | 19 | 0 | 19 | 0 | 19 | 210 | 0 | 0 |
| Michigan | 141 | 0 | 140 | 6 | 141 | 288 | 6 | 12 |
| Minnesota | 13 | 0 | 13 | 1 | 13 | 142 | 1 | 11 |
| Missouri | 75 | 0 | 70 | 4 | 75 | 238 | 4 | 13 |
| Nebraska | 8 | 0 | 8 | 0 | 8 | 187 | 0 | 0 |
| North Dakota | 1 | 0 | 1 | 0 | 1 | 75 | 0 | 0 |
| Ohio | 122 | 1 | 121 | 3 | 122 | 271 | 3 | 7 |
| South Dakota | 4 | 0 | 4 | 0 | 4 | 120 | 0 | 0 |
| Wisconsin | 31 | 2 | 29 | 2 | 31 | 137 | 2 | 9 |
| South | 1,565 | 50 | 1,483 | 79 | 1,568 | 259 | 79 | 13 |
| Alabama | 75 | 0 | 66 | 2 | 75 | 270 | 2 | 7 |
| Arkansas | 58 | 3 | 38 | 3 | 58 | 431 | 3 | 22 |
| Delaware | 22 | 0 | 19 | 3 | 22 | 306 | 3 | 42 |
| Florida | 248 | 14 | 244 | 17 | 248 | 283 | 17 | 19 |
| Georgia | 128 | 10 | 122 | 10 | 128 | 268 | 10 | 21 |
| Kentucky | 41 | 1 | 43 | 1 | 43 | 228 | 1 | 5 |
| Louisiana | 103 | / | 89 | 7 | 103 | 276 | 7 | 19 |
| Maryland | 57 | / | 57 | 12 | 57 | 245 | 12 | 52 |
| Mississippi | 65 | 2 | 64 | 2 | 65 | 312 | 2 | 10 |
| North Carolina | 73 | 1 | 70 | 1 | 73 | 201 | 1 | 3 |
| Oklahoma | 79 | 2 | 75 | 2 | 79 | 333 | 2 | 8 |
| South Carolina | 76 | 3 | 75 | 3 | 76 | 318 | 3 | 13 |
| Tennessee | 75 | 0 | 76 | 2 | 76 | 290 | 2 | 8 |
| Texas | 371 | 10 | 351 | 10 | 371 | 217 | 10 | 6 |
| Virginia | 84 | 3 | 84 | 3 | 84 | 236 | 3 | 8 |
| West Virginia | 10 | 1 | 10 | 1 | 10 | 190 | 1 | 19 |
| West | 709 | 20 | 680 | 21 | 714 | 241 | 25 | 8 |
| Alaska | 0 | / | 4 | 0 | 4 | 86 | 0 | 0 |
| Arizona | 77 | 1 | 78 | 1 | 78 | 239 | 1 | 3 |
| California | 374 | 9 | 366 | 14 | 374 | 225 | 14 | 8 |
| Colorado | 65 | 4 | 54 | 3 | 65 | 312 | 4 | 19 |
| Hawaii | 8 | 0 | 8 | 0 | 8 | 132 | 0 | 0 |
| Idaho | 16 | / | 13 | 0 | 16 | 261 | 0 | 0 |
| Montana | 14 | 0 | 12 | 0 | 14 | 416 | 0 | 0 |
| Nevada | 33 | 0 | 29 | 0 | 33 | 285 | 0 | 0 |
| New Mexico | 19 | 0 | 18 | 0 | 19 | 288 | 0 | 0 |
| Oregon | 41 | 2 | 37 | 2 | 41 | 308 | 2 | 15 |
| Utah | 8 | 0 | 8 | 0 | 8 | 133 | 0 | 0 |
| Washington | 46 | 4 | 45 | 1 | 46 | 276 | 4 | 24 |
| Wyoming | 8 | 0 | 8 | 0 | 8 | 395 | 0 | 0 |

/Not reported.

^aBased on individual reports submitted to the Deaths in Custody Reporting Program (DCRP).

^bData from the National Prisoners Statistics (NPS-1) and DCRP were combined to provide the number of total deaths in each State. The most complete data for each State were used to calculate a total for each State.

Appendix table 5. Circumstances under which inmates were tested for the antibody to the human immunodeficiency virus (HIV), by jurisdiction, 2005

| Jurisdiction | All inmates | | | High-risk group | Inmate request | Clinical indication | Involvement in incident | Random sample | Court order | Other |
|------------------|-------------|------------|--------------|-----------------|----------------|---------------------|-------------------------|---------------|-------------|-------|
| | Entering | In custody | Upon release | | | | | | | |
| Total | | | | ■ | ■ | ■ | ■ | | ■ | |
| Federal system* | | | | ■ | ■ | ■ | ■ | | ■ | |
| Northeast | | | | | | | | | | |
| Connecticut | | | | ■ | ■ | ■ | ■ | | ■ | |
| Maine | | | | | ■ | ■ | | | ■ | |
| Massachusetts | | | | | ■ | | | | | |
| New Hampshire | ■ | | | | | | | | | |
| New Jersey | | | | | ■ | ■ | | | ■ | |
| New York | | | | ■ | ■ | ■ | ■ | ■ | ■ | |
| Pennsylvania | | | | | ■ | ■ | ■ | | ■ | |
| Rhode Island | ■ | | | | ■ | ■ | ■ | | ■ | |
| Vermont | | | | | ■ | ■ | ■ | | | |
| Midwest | | | | | | | | | | |
| Illinois | | | | ■ | ■ | ■ | ■ | | ■ | |
| Indiana | ■ | ■ | | ■ | ■ | ■ | | | ■ | |
| Iowa | ■ | | | | | ■ | ■ | | | |
| Kansas | | | | ■ | ■ | ■ | ■ | | ■ | |
| Michigan | ■ | | | | ■ | | ■ | | ■ | |
| Minnesota | | | | | ■ | ■ | ■ | | ■ | |
| Missouri | ■ | | ■ | | ■ | ■ | ■ | | ■ | |
| Nebraska | ■ | | | | ■ | ■ | ■ | | ■ | |
| North Dakota | ■ | | | | ■ | ■ | ■ | | ■ | |
| Ohio | ■ | | | | ■ | ■ | ■ | | ■ | |
| South Dakota | | | | | ■ | ■ | ■ | | ■ | |
| Wisconsin | | | | | ■ | ■ | ■ | | ■ | ■ |
| South | | | | | | | | | | |
| Alabama | ■ | ■ | ■ | | | ■ | ■ | | ■ | |
| Arkansas | ■ | ■ | | ■ | ■ | ■ | ■ | ■ | ■ | |
| Delaware | | | | | ■ | ■ | ■ | | ■ | |
| Florida | | | ■ | | ■ | ■ | ■ | | | ■ |
| Georgia | ■ | | | | ■ | ■ | ■ | | ■ | |
| Kentucky | | | | ■ | | ■ | ■ | | ■ | |
| Louisiana | | | | | ■ | ■ | ■ | | ■ | |
| Maryland | | | | | ■ | ■ | ■ | | ■ | ■ |
| Mississippi | ■ | | | ■ | ■ | ■ | ■ | | ■ | |
| North Carolina | | | | | ■ | ■ | ■ | | ■ | |
| Oklahoma | ■ | | | | ■ | ■ | ■ | | ■ | ■ |
| South Carolina | ■ | | | | ■ | ■ | ■ | | ■ | ■ |
| Tennessee | | | | | ■ | ■ | ■ | | ■ | |
| Texas | | | ■ | ■ | ■ | ■ | ■ | | ■ | |
| Virginia | | | | | ■ | ■ | ■ | | | |
| West Virginia | | | | | ■ | ■ | ■ | | | |
| West | | | | | | | | | | |
| Alaska | / | / | / | / | / | / | / | / | / | / |
| Arizona | | | | | ■ | ■ | | | ■ | |
| California | | | | | ■ | ■ | | | ■ | |
| Colorado | ■ | | | | ■ | ■ | ■ | | ■ | |
| Hawaii | | | | | ■ | ■ | ■ | | ■ | |
| Idaho | ■ | ■ | | ■ | ■ | ■ | ■ | | ■ | |
| Montana | | | | | ■ | ■ | ■ | | ■ | |
| Nevada | | ■ | ■ | | ■ | ■ | ■ | ■ | ■ | |
| New Mexico | | | | | ■ | ■ | | | ■ | |
| Oregon | | | | ■ | ■ | ■ | ■ | ■ | ■ | |
| Utah | ■ | | | ■ | | ■ | ■ | | | |
| Washington | | | | ■ | ■ | ■ | ■ | | ■ | |
| Wyoming | ■ | | | | ■ | ■ | | | | |

/Not reported.

*The Bureau of Prisons tests a random sample of inmates in alternate years.

Appendix table 6. Inmate deaths in Federal prisons, by cause, 2004 and 2005

| Cause of death | Deaths of Federal inmates reported in the NPS-1 | | | |
|--------------------------------|---|------|---------------------------|------|
| | Number | | Rate per 100,000 inmates* | |
| | 2005 | 2004 | 2005 | 2004 |
| Total | 388 | 333 | 210 | 186 |
| Natural causes other than AIDS | 327 | 295 | 177 | 165 |
| AIDS | 27 | 18 | 15 | 10 |
| Suicide | 13 | 11 | 7 | 6 |
| Accident | 6 | 6 | 3 | 3 |
| By another person | 15 | 3 | 8 | 2 |

*Detail may not add to total due to rounding.

Appendix table 7. Standard error estimates for Survey of Inmates in State Correctional Facilities, 1997 and 2004

| Characteristic | Estimated standard error of the percent of inmates who were HIV positive | |
|-----------------------------|--|-------|
| | 2004 | 1997 |
| Gender | | |
| Male | 0.20% | 0.20% |
| Female | 0.35 | 0.44 |
| Race/Hispanic origin | | |
| White* | 0.30% | 0.29% |
| Black or African American* | 0.34 | 0.37 |
| Hispanic or Latino | 0.47 | 0.58 |
| Age | | |
| 24 or younger | 0.23% | 0.20% |
| 25-34 | 0.25 | 0.30 |
| 35-44 | 0.46 | 0.40 |
| 45 or older | 0.61 | 0.60 |
| Current offense | | |
| Violent | 0.27% | 0.25% |
| Drug | 0.48 | 0.46 |
| Property | 0.60 | 0.40 |
| Public-order | 0.46 | 0.55 |
| Prior drug use | | |
| Never | 0.46% | 0.41% |
| Ever | 0.23 | 0.21 |
| In month before arrest | 0.29 | 0.27 |
| Used needle to inject drugs | 0.62 | 0.57 |
| Shared a needle | 1.27 | 1.09 |

*Not Hispanic or Latino.